

ORAL PRESENTATION

Open Access

Quantum-mechanics-based molecular interaction fields for 3D-QSAR

Ahmed ElKerdawy^{1*}, Stefan Güssregen², Hans Matter², Matthias Hennemann^{1,3}, Timothy Clark^{1,3,4}

From 9th German Conference on Chemoinformatics
Fulda, Germany. 10-12 November 2013

Computer-aided drug design (CADD) shift toward using quantum-mechanics (QM)-based approaches is not only the result of the ever growing computational power but also due to the need for more accurate and more informative approaches to describe molecular properties and binding characteristics than the currently available ones. QM-approaches do not suffer from the limitations inherent to the ball-and-spring description and the fixed atom-centered charge approximation in the classical

force fields mostly used by CADD methods.[1] We introduce a protocol for shifting 3D-QSAR, one of the most widely used ligand-based drug design approaches, through using QM-based molecular interaction fields (MIFs) which are the electron density (ρ), hydrogen bond donor field (HDF), hydrogen bond acceptor field (HAF) and molecular lipophilicity potential (MLP) to overcome the limitations of the current force-field-based MIFs (FF-MIFs). The average performance of the




Figure 1 Performance of the different 3D-QSAR approaches in the external validation.

* Correspondence: Ahmed.ElKerdawy@fau.de

¹Computer-Chemie-Centrum, Friedrich-Alexander-Universität Erlangen-Nürnberg, Nögelsbachstraße 25, 91052 Erlangen, Germany
Full list of author information is available at the end of the article

QM-MIFs (QMFA) models for nine data sets was found to be better than that of the conventional FF-MIFs models. In the individual data sets, the QMFA models always perform better than, or as well as, the conventional approaches. It is particularly encouraging that the relative performance of the QMFA models improves in the external validation (Figure 1).

Authors' details

¹Computer-Chemie-Centrum, Friedrich-Alexander-Universität Erlangen-Nürnberg, Nögelsbachstraße 25, 91052 Erlangen, Germany. ²Sanofi-Aventis Deutschland GmbH, R&D, LGCR, Structure, Design and Informatics, Building G 878, 65926, Frankfurt am Main, Germany. ³Interdisciplinary Center for Molecular Materials, Friedrich-Alexander-Universität Erlangen-Nürnberg, Nögelsbachstraße 49, 91052 Erlangen, Germany. ⁴Centre for Molecular Design, University of Portsmouth, King Henry Building, Portsmouth PO1 2DY, UK.

Published: 11 March 2014

Reference

1. Zhou T, Huang DZ, Caflisch A: **Quantum Mechanical Methods for Drug Design.** *Curr Top Med Chem* 2010, **10**(1):33-45.

doi:10.1186/1758-2946-6-S1-O10

Cite this article as: ElKerdawy et al.: Quantum-mechanics-based molecular interaction fields for 3D-QSAR. *Journal of Cheminformatics* 2014 **6**(Suppl 1):O10.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

"Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge."

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>



ChemistryCentral